


2839
2840
2841
2842
2843
2844
2845
2846
2848
2849
2850
2852
2853
2854
2855
2856
2857
2859
2860

K. Nafstad Ingebrigtsen. Distribution and transplacental transfer of paraquat in rats and guinea-pigs. General Pharmacology: The Vascular System. 1984. 15:201-204
J. P. Fouquet Morin. Development of a new in vitro system for continuous in vitro exposure of lung tissue to complex atmospheres: Application to diesel exhaust toxicology. Cell Biology and Toxicology. 1999. 15:143-152
B. H. Wu Kang. Detrimental effects of exogenous nitric oxide in paraquat-induced lung injury. Free Radical Biology and Medicine. 2001. 31:S70-S70
D. T. Wang Kong. Determination of Anti-Tumor Agent TM208, 4-Methyl-piperazine-1-carbodithioc Acid 3-Cyano-3,3-diphenylpropyl Ester Hydrochloride, in Rats by LC. Chromatographia. 2009. 70:1721-1725
P. I. Friedman Karl. Competition between paraquat and putrescine for accumulation by rat lung slices. Toxicology. 1983. 26:317-323
Basim A. Maxwell Dubaybo. Compensatory Growth in Fibrotic Lung Injury. The American Journal of the Medical Sciences. 1991. 302:19-22
H. F. Takezawa Lam. A comparison of the effects of paraquat and diquat on lung compliance, lung volumes and single breath diffusing capacity in the rat. Toxicology. 1980. 18:111-123
J. R. Acuff Dunbar. CO-ADMINISTRATION OF PARAQUAT AND PUTRESCINE TO RATS VIA MINIOSMOTIC PUMP - EFFECTS ON LUNG GLUTATHIONE ANTIOXIDANT SYSTEM AND PQ CONTENT. Federation Proceedings. 1985. 44:1024-1024
A. Verma Mehta. Chlorpyrifos induced alterations in the levels of hydrogen peroxide, nitrate and nitrite in rat brain and liver. Pesticide Biochemistry and Physiology. 2009. 94:55-59
G. T. Pilarski Chaneva. CHANGES OF PROLINE CONTENT IN A CYANOBACTERIUM UNDER OXIDATIVE STRESS. Oxidation Communications. 2011. 34:439-445
In Youb Yoon Chang. The changes of calretinin immunoreactivity in paraquat-induced nephrotoxic rats. Acta Histochemica. 2012. 114:836-841
Watanabe Shoichi Satoru. Change of polyamine level in various tissues of male rats intoxicated with paraquat. Toxicology Letters. 1993. 69:163-170
J. B. Hill Atkinson. Centrilobular endothelial cell injury by diquat in the selenium-deficient rat liver. Laboratory Investigation. 2001. 81:193-200
K. Hasegawa Yamanaka. CELLULAR-RESPONSE TO OXIDATIVE DAMAGE IN LUNG INDUCED BY THE ADMINISTRATION OF DIMETHYLARSINIC ACID, A MAJOR METABOLITE OF INORGANIC ARSENICS, IN MICE. Toxicology and Applied Pharmacology. 1991. 108:205-213
Y. X. Cheng Fui. Cellular glutathione peroxidase protects mice against lethal oxidative stress induced by various doses of diquat. Proceedings of the Society for Experimental Biology and Medicine. 1999. 222:164-169
Hossein Sanaei-Zadeh. Can pirfenidone prevent paraquat-induced pulmonary fibrosis?—A hypothesis. Tzu Chi Medical Journal. 2012. 24:223
D. Bagchi Bagchi. Cadmium-induced excretion of urinary lipid metabolites, DNA damage, glutathione depletion, and hepatic lipid peroxidation in Sprague-Dawley rats. Biological Trace Element Research. 1996. 52:143-154
V. C. Nwanze Nwabisi. Biochemical studies on the toxicity of 1,1'-dimethyl-4,4'-bipyridylium dichloride in the rat. Comparative Biochemistry and Physiology C-Pharmacology Toxicology & Endocrinology. 1997. 117:103-109
R. A. Ingebrigtsen Andersen. Biochemical characteristics of rat superoxide dismutase and the effect caused by paraquat injection on the enzyme activity in various tissues. General Pharmacology: The Vascular System. 1984. 15:205-210

1. 1. Radiolabelled paraquat (bis-N-(14C)methyl-4,4'-bipyridylium chloride) was administered intravenously to pregnant
The purpose of this study was the development of a new incubation system that can allow continuous exposure of lung t
A new, specific and sensitive high performance liquid chromatography analytical procedure was developed and validated
Slices from rat lungs actively accumulate both paraquat and putrescine, while endogenous polyamines inhibit the accum
ABSTRACT We induced severe left-sided lung fibrosis by the unilateral endobronchial instillation of paraquat (1.0 mg/kg)
Paraquat intoxication in its initial stage is characterized histologically in the lungs by atelectasis, hyaline membrane form
Organophosphate pesticides are among the most widely used synthetic chemicals for controlling a wide variety of pests.
Changes of free proline content in <i>Plectonema boryanum</i> cells after cultivation at oxidative stress inducing conditions we
Calcium-binding proteins are present in the kidneys: calbindin D-28k in the distal tubules and calretinin in the proximal tu
The effects of paraquat (PQ) on the contents of putrescine (PUT), spermidine (SPD) and spermine (SPM) in the various tiss
Low doses of diquat cause massive liver necrosis and death of selenium-deficient rats within a few hours. Protection aga
This study was to determine if cellular glutathione peroxidase (GPX1) protects against acute oxidative stress induced by c
Recent studies have described lipid peroxidation to be an early and sensitive consequence of cadmium exposure, and fre
The effect of intraperitoneal administration of lethal dose (50 mg/kg) of paraquat on the microsomal cysteine levels in th
1. 1. Rat superoxide dismutase (SOD) prepared from the rat was not found by sedimentation in sucrose gradients or isole

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



2861
2862
2863
2865
2866
2867
2868
2869
2870
2871
2872
2874
2875
2876
2877
2878
2879
2880
2881
2882

Shri N. Lunsman Giri. Binding of [methyl-3H]paraquat to rat, rabbit, hamster, mouse and guinea pig lung proteins, in vitro. <i>Toxicology Letters</i> . 1981. 9:93-100
R. D. Goncalves Novaes. Bark extract of <i>Bathysa cuspidata</i> attenuates extra-pulmonary acute lung injury induced by paraquat and reduces mortality in rats. <i>International Journal of Experimental Pathology</i> . 2012. 93:225-233
Takahiko Taguchi Sato. Augmentation of allergic reactions by several pesticides. <i>Toxicology</i> . 1998. 126:41-53
M. D. Lison Schweich. ASSESSMENT OF LIPID-PEROXIDATION ASSOCIATED WITH LUNG DAMAGE-INDUCED BY OXIDATIVE STRESS - IN-VIVO AND IN-VITRO STUDIES. <i>Biochemical Pharmacology</i> . 1994. 47:1395-1400
Timothy M. Montgomery Sullivan. Ascorbic acid nutritional status does not affect the biochemical response to paraquat. <i>Fundamental and Applied Toxicology</i> . 1984. 4:754-759
S. Suncica Ivanovic. The application of adsorbent bentonite in oxidative stress induced by paraquat. <i>Acta Veterinaria-Beograd</i> . 2007. 57:329-340
Satoshi Iwasaka Hagiwara. An antisense oligonucleotide to HSP47 inhibits paraquat-induced pulmonary fibrosis in rats. <i>Toxicology</i> . 2007. 236:199-207
B. J. Day. Antioxidants as potential therapeutics for lung fibrosis. <i>Antioxidants & Redox Signaling</i> . 2008. 10:355-370
T. W. Wolfgang Petry. ANTIOXIDANT-DEPENDENT INHIBITION OF DIQUAT-INDUCED TOXICITY INVIVO. <i>Toxicology</i> . 1992. 74:33-43
Y. Kim Kim. Antioxidant and anti-inflammatory effects of selenium in oral buccal mucosa and small intestinal mucosa during intestinal ischemia-reperfusion injury. <i>Journal of Inflammation-London</i> . 2014. 11:#pages#
A. F. Hasan Mohamed. Antioxiclant and hepatoprotective effects of <i>Eucalyptus maculata</i> . <i>Medical Science Monitor</i> . 2005. 11:BR426-BR431
Hassan Mehrabi Malekinejad. Antifibrotic effect of atorvastatin on paraquat-induced pulmonary fibrosis: Role of PPAR γ receptors. <i>European Journal of Pharmacology</i> . 2013. 720:294-302
J. T. Hardwick Skamarauskas. Adenosine accumulation as an indicator of cell-specific toxicity in rat lung slices. <i>Toxicology in Vitro</i> . 1995. 9:757-763
Edward A. Ishmael Lock. The acute toxic effects of paraquat and diquat on the rat kidney. <i>Toxicology and Applied Pharmacology</i> . 1979. 50:67-76
Lewis L. Pratt Smith. The accumulation of putrescine and paraquat into lung slices taken from BHT treated mice. <i>Toxicology</i> . 1983. 27:13-Jan
S. Smith Brooke-Taylor. The accumulation of polyamines and paraquat by human peripheral lung. <i>Biochemical Pharmacology</i> . 1983. 32:717-720
G. Honarjoo Nasiri. 392 C0-Administration of captopril or niacin with paraquat in isolated rat lung perfusion. <i>Toxicology Letters</i> . 2003. 144, Supplement <HT>1</HT>:s106-s107
M. Oshida J. Orino K. Watanabe K. Higuchi. Wheat Bran Protects Fischer-344 Rats from Diquat-Induced Oxidative Stress by Activating Antioxidant System: Selenium as an Antioxidant. <i>Bioscience Biotechnology and Biochemistry</i> . 2011. 75:496-499
O. Diaz P. Agius M. F. Valpuesta V. Monza J. Borsani. Water stress generates an oxidative stress through the induction of a specific Cu/Zn superoxide dismutase in <i>Lotus corniculatus</i> leaves. <i>Plant Science</i> . 2001. 161:757-763
A. Demirevska K. Gesheva E. Gechev T. Edreva. UP-REGULATION OF ROS SCAVENGING ENZYMES IN atr ARABIDOPSIS THALIANA MUTANTS WITH COMPROMISED PROGRAMMED CELL DEATH. <i>Comptes Rendus De L Academie Bulgare Des Sciences</i> . 2011. 64:1131-1136

When incubated with the rat lung slices [methyl-3H] paraquat was found to bind covalently to acid-insoluble proteins. Th
This study investigated the effect of the bark extract of <i>Bathysa cuspidata</i> on paraquat (PQ)-induced extra-pulmonary ac
The augmentative effects of several pesticides on histamine release from mast cells of rats that had been sensitized pass
The lung thiobarbituric acid-reactive substances (TBA-RS) content and the amount of ethane exhaled, two potential mark
Guinea pigs were subjected to dietary control of ascorbic acid intake to produce deficient, normal, and supplemented tis
The protective effect of bentonite (natural and synthetic) in oxidative stress induced by paraquat was studied on 32 adult
The most common cause of death from poisoning by the widely used, but highly toxic herbicide paraquat is respiratory fa
Interstitial lung disease encompasses a large group of chronic lung disorders associated with excessive tissue remodeling
The abilities of two experimental antioxidants (U-74006F and U-78517G), as well as the model antioxidant, diphenyl-p-ph
Background: The aim of this study were to investigate whether selenium treatment attenuates lipid peroxidation and do
Background: Since phenolic compounds have been reported as effective antioxidants, this study was designed to assess t
This study was carried out to highlight the role of PPAR? in the paraquat (PQ)-induced pulmonary fibrosis. Forty-two mal
We have investigated the potential of adenosine uptake as a marker of chemically induced, cell-selective pulmonary inju
The bipyridyls, paraquat and diquat, cause mild renal tubular damage in the rat. A marked diuresis, albuminuria, glucosu
Putrescine and paraquat have been shown to accumulate in slices of mouse lung by a process which obeys saturation kin
Wheat bran had a protective effect against diquat toxicity in rats fed a purified diet (PD). We studied the effects of wheat
Drought brings about different biochemical responses in plants in order to minimize its deleterious effects. Drought indu
This work describes the alteration in the activities of a range of ROS-scavenging enzymes and nonenzymatic antioxidants

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant